



Detection of polytreponemal infection in three cases of porcine ulcerative stomatitis by Fluorescent in situ hybridization

Jensen, Tim Kåre; Strijkstra, G.T.; Gruys, E.; Baumgärtner, W.; Schou, Kirstine Klitgaard; Boye, Mette

Publication date:
2014

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Jensen, T. K., Strijkstra, G. T., Gruys, E., Baumgärtner, W., Schou, K. K., & Boye, M. (2014). *Detection of polytreponemal infection in three cases of porcine ulcerative stomatitis by Fluorescent in situ hybridization*. Abstract from Cutting Edge Pathology, Berlin, Germany.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Your Abstract

Detection of polytreponemal infection in three cases of porcine ulcerative stomatitis by Fluorescent in situ hybridization

T. K. Jensen^{*}, G.T. Strijkstra[†], E. Gruys[‡], W. Baumgärtner^{}, K. Klitgaard^{*} and M. Boye^{*}**

^{*}National Veterinary Institute, Technical University of Denmark, Denmark, [†]Tierärztliche Gemeinschaftspraxis BSB, Bersenbrück, Germany, [‡]Veterinary Extension Services, Driebergen, Netherlands, ^{**}Department of Pathology, University of Veterinary Medicine, Hannover, tije@vet.dtu.dk

Introduction: Ulcerative and fibrino-necrotizing stomatitis is occasionally reported in pigs. The etiology is unknown but spirochetes are observed in silver stained sections, however, the spirochetes have never been identified. The aim of this study was to report three cases of fibrino-necrotizing stomatitis associated with polytreponemal infection.

Materials and Methods: Tissue samples from the oral cavity of three sows euthanized due to ulcerative stomatitis, ulcerative vaginitis and Mortellaro-like alterations of the feet were fixed in formalin, paraffin embedded and sectioned for H&E and serial fluorescent in situ hybridization (FISH) for identification of spirochetes with genus and species specific oligonucleotide probes targeting 16S or 23S ribosomal RNA. The presence of *Fusobacterium necrophorum* and *Domain Bacteria* was additionally detected by FISH.

Results: Diffuse fibrino-necrotizing infection affecting the epidermis and dermis of the entire oral cavity was found in all sows. By FISH all lesions showed severe bacterial infiltration with a strong predominance of spirochetes. Additional hybridization revealed that the spirochetes all belonged to genus *Treponema*. Neither, *Brachyspira*, *Borrelia*, nor *Leptospira* organisms were detected. Further identification of the spirochetes showed intermingled infection with *Treponema pedis*, *Treponema putidum* and up to eight other *Treponema* species/phylotypes. Low numbers of *F. necrophorum* was also detected but only the treponemes were infiltrating deep into the lesions to the borderline between damaged and vital tissue.

Discussion: Using a panel of genus specific probes this is the first study identifying the involvement of *Treponema spp.* in porcine ulcerative stomatitis. All three sows revealed polytreponemal infections with *Treponema* species/phylotypes commonly associated with bovine digital dermatitis and human periodontal disease.